

Symmetry breaking and structural distortions in charged XH_4 ($\text{X} = \text{C}, \text{Si}, \text{Ge}, \text{Sn}$ and Pb) molecules

D. Balamurgan, Manoj. K. Harbola and R. Prasad

Department of Physics, Indian Institute of Technology, Kanpur 208016, India

Abstract

We have investigated the ground state structures of neutral and charged XH_4 ($\text{X}=\text{C}, \text{Si}, \text{Ge}, \text{Sn}$ and Pb) molecules using the first-principles electronic structure methods. The structure of positively charged molecules for $\text{X} = \text{Si}, \text{Ge}, \text{Sn}$ and Pb is characterized by a severe distortion from tetrahedral structure and an unusual H-H bond while the negatively charged molecules get distorted by pushing two hydrogen atoms away from each other. However, CH_4^+ and CH_4^- are exceptions to this behavior. We provide an insight into the symmetry breaking mechanism and unusual H-H bonding using simple electrostatic arguments based on the unequal charge distribution on H atoms. Those charged molecules having unequal charge distribution on H atoms get distorted due to different electrostatic forces between the atoms. We show that the directionality and occupation of the highest occupied molecular orbital(HOMO) play an important role in creating charge asymmetry in these molecules.

PACS numbers: 73.22.-f, 36.40.Mr, 36.40.Wa, 36.40.Qv

I. INTRODUCTION

Although neutral XH_4 ($\text{X} = \text{C}, \text{Si}, \text{Ge}, \text{Sn}$ and Pb) molecules have tetrahedral symmetry, charging these molecules breaks this symmetry. The symmetry breaking in general is associated with a large structural distortion and in many cases unusual H-H bonding arrangement. Some of these clusters have been investigated earlier and the symmetry breaking has been attributed to the Jahn-Teller effect. In the Jahn-Teller effect[1, 2, 3], electrons occupying a degenerate level favor a low symmetric structure by lifting the degeneracy. For example, the highest occupied molecular orbital(HOMO) of tetrahedral SiH_4 is triply degenerate. When an electron is removed from SiH_4 , its structure gets distorted drastically due to the Jahn-Teller effect[4, 5, 6, 7, 8].

The structure of SiH_4^+ has been investigated by a large number of workers[4, 5, 6, 7, 8] and it took almost a decade to settle its ground state structure. Now it is generally agreed that the ground state structure of SiH_4^+ has C_s symmetry. Similarly, the structure of CH_4^+ has been studied extensively[8, 9, 10, 11, 12, 13, 14, 15, 16], though there is still ambiguity regarding its ground state structure. The calculations done by the density functional theory(DFT) methods show that the ground state structure of CH_4^+ has D_{2d} symmetry[8, 9], while the Hartree-Fock(HF) and post HF calculations show the ground state structure as C_{2v} [9, 16]. The lowering of tetrahedral symmetry of CH_4^+ and SiH_4^+ has been attributed to the Jahn-Teller effect. However, the structure of CH_4^+ is quite different from that of SiH_4^+ . It is difficult to understand the difference between the CH_4^+ and SiH_4^+ distortions only from the Jahn-Teller effect. This is because the Jahn-Teller theorem only suggests the existence of distortion and says nothing regarding the nature and magnitude of the distortion.

In this paper we focus on the ground state structures of charged XH_4 molecules using the first-principles electronic structure methods and provide an insight into the symmetry breaking mechanism. We find that upon charging most of the molecules undergo large structural distortion which can be attributed to the Jahn-Teller effect. Further, the nature of distortions of positively charged molecules is different from that of negatively charged molecules. Interestingly, most of the positively charged molecules have an unusual H-H bond and appear like transition states of unstable XH_2^+ and H_2 complex. We try to understand the distortions, unusual H-H bonding and stability of these molecules. Vibrational analysis shows that these molecules are not transition states. The calculated fragmentation energy

indicates that the stability of XH_4^+ molecules decreases from CH_4^+ to PbH_4^+ .

Using various exchange-correlation functionals such as local density approximation(LDA), local spin density approximation(LSDA), gradient corrected approximation(GGA) and spin-polarized GGA(GGA-SP) we have investigated the effect of different occupations in charged tetrahedral XH_4 molecules. Our calculations show that the configuration corresponding to integer occupancy and asymmetrical charge density leads to the lowest energy. We provide simple arguments in terms of charge asymmetry and electrostatic interaction between atoms which explain the nature of distortions in charged XH_4 molecules. For example, we are able to explain why the nature of distortion of SiH_4^+ is different from CH_4^+ and why the positively charged molecules distort differently from the negatively charged molecules. We study the structural distortions of SiH_4^+ in detail using conjugate gradient method and show that the evolution of the structure is governed by the electrostatic interaction between the atoms. Our investigations reveal that the distribution of charges on H atoms plays an important role in distorting the charged molecules. If the charges on H atoms are not equal, the structure gets distorted due to different electrostatic forces on the atoms. We also discuss how the charge asymmetry is created in these molecules by the directionality and occupation of the HOMO.

The plan of the paper is as follows. In section **II** we give computational details of the present work. Section **III** contains results and discussion and finally in section **IV** we give our conclusions.

II. COMPUTATIONAL DETAILS

We have investigated the structure of neutral and charged XH_4 using Vienna *ab-initio* simulation package(VASP)[17] and Gaussian98[18] package. We have cross checked our calculations using both packages and find that the results obtained by both methods are in good agreement. A number of calculations were done using the unrestricted Hartree-Fock(UHF) and DFT methods using the Gaussian98 package. In the DFT calculations we have used several exchange-correlation functionals such as Slater's exchange with the local correlation functional of Perdew(SPL), Becke's exchange which includes gradient density with gradient corrected correlation functional(BPW91) and Becke's 3 parameter hybrid functional(B3LYP)[18, 19]. In these calculations we have employed 6-311g** basis set for

molecules having C and Si atoms and LanL2DZ basis set with effective core potential for other molecules. The plane wave ultrasoft pseudopotential calculations[17] were done with the local density approximation(LDA)[20, 21] employing a simple cubic supercell of $20\text{\AA}$. The number of plane waves used in the calculation is set by energy cut-off(Ecut). We have used 287 eV Ecut for neutral and charged CH_4 molecules and 150 eV Ecut for other molecules. The use of supercell technique for charged systems causes electrostatic interaction within the images of supercells[22, 23]. Calculation of charged systems are efficiently handled in VASP by applying a background charge to maintain the charge neutrality and by adding dipole and quadrupole moment corrections[17]. The structure and ionization potential of SiH_4^+ predicted by using this procedure are in close agreement with those of earlier calculations based on the HF, MP2 and LSDA methods[6, 8].

III. RESULTS AND DISCUSSION

A. Ground state structures

Fig. 1 shows the ground state structures of XH_4 molecules under neutral and charged states using the Gaussian98 package with BPW91 functional. We first discuss the ground state structures of the positively charged molecules. We see from the figure that the tetrahedral symmetry is broken when XH_4 molecules are ionized. The structure of CH_4^+ has D_{2d} symmetry, while other positively charged molecules have C_s symmetry. We have investigated various possible symmetries of XH_4^+ molecules and our results are summarized in Table. I. We see from the table that C_{2v} and D_{2d} structures of CH_4^+ are very close in energy and differ only by 0.11 eV. The other positively charged molecules also have C_{2v} as the next higher energy structure. We could not find local minima of C_{3v} and C_s symmetries for CH_4^+ and C_{2v} symmetry for SiH_4^+ at the BPW91 level. These missing structures of CH_4^+ and SiH_4^+ molecules in our calculations have been studied earlier using different exchange-correlation potentials and different methods[4, 5, 6, 9, 15, 16] which show that these are not the ground state structures.

It is interesting to note that in the C_s symmetry, the positively charged molecules have a structure which is severely distorted from the tetrahedral structure. Furthermore, there is an unusual bonding between two hydrogen atoms. This bonding is confirmed by the charge

density calculation as shown in Fig. 2 which shows large electron density between the two hydrogen atoms of SiH_4^+ . To ascertain that this is not an artifact of the BPW91 calculation, we have re-optimized the ground state structures using the HF and DFT methods with different exchange correlation functionals such as SPL and B3LYP. The H-H bond lengths obtained from various methods are essentially the same. This kind of over-coordination of hydrogen has also been observed in CH_5^+ [24, 25], SiH_5^+ [26] and small hydrogenated silicon clusters[27]. In Fig. 3, we have shown the total energy curve of SiH_4^+ , H_2 and H_2^+ as a function of H-H distance. The total energy of SiH_4^+ is calculated by keeping Si-H bond length fixed but varying H-H bond distance. From the figure we see that the curvature of SiH_4^+ curve is smaller than H_2 and larger than H_2^+ curves. This shows H-H bond in SiH_4^+ is stronger than H_2^+ but weaker than H_2 . Vibrational analysis also indicates that the H-H bond is sufficiently strong in these molecules. We see from Table. II that the force constant for H-H stretch mode of H-H pair is smaller than H_2 and larger than H_2^+ which implies that the strength of H-H bond is weaker than H_2 and stronger than H_2^+ .

For the ground state structures of the negatively charged molecules, we note from Fig. 1 that the distortions of negatively charged molecules are different from those of positively charged molecules. XH_4^- molecules have C_{2v} symmetry except CH_4^- which has tetrahedral symmetry. The two hydrogen atoms of XH_4^- molecules are pushed away from each other which is in contrast to the H-H bond of XH_4^+ molecules with C_s symmetry.

B. Stability

One may argue that the structure of XH_4^+ molecules with C_s symmetry appears more like XH_2^+ and H_2 complex and hence unstable. Frey and Davidson[4] studied SiH_4^+ using CI singles root calculation and showed that C_s structure is more stable than C_{2v} against fragmentation into SiH_2^+ and H_2 . We have calculated the fragmentation energy which is the energy required to fragment a XH_4^+ molecule into its binary products and has been calculated by taking the total energy difference between the XH_4^+ and the possible binary products. Of all the possible channels, the fragmentation of XH_4^+ into XH_2^+ and H_2 requires least energy. We give the fragmentation energies of XH_4^+ molecules for this process in Table. III. We see from the table that the energy required for the fragmentation of XH_4^+ into XH_2^+ and H_2 decreases for $\text{X}=\text{C}$ to Pb . This means that the stability of XH_4^+ molecules decreases

from C to Pb molecules.

To examine the stability further we have done vibrational analysis of these molecules. In Table. IV we present the vibrational frequencies of charged and neutral XH_4 molecules. Our calculated vibrational frequencies 880, 950, 2175 and 2192cm^{-1} of SiH_4 are in good agreement with those of experimentally obtained values[28] of 914, 953, 2189 and 2267cm^{-1} . We see from the table that PbH_4^+ has a negative frequency and other molecules have only positive frequencies. Further, we see that one frequency is very low for C_s symmetric XH_4^+ molecules and decreases from SiH_4^+ to GeH_4^+ and become negative for PbH_4^+ .

The positive frequencies of a molecule indicate that the structure corresponds to an energy minimum. If one frequency is negative, the structure may be a transition state of the molecule[19]. Since the frequency calculation is highly dependent on the method, the small negative frequency of PbH_4^+ may be an artifact of the method. To clarify this we have done vibrational analysis on re-optimized structures of C_s symmetric XH_4^+ molecules using various methods. Calculation with the SPL exchange-correlation functional shows that one frequency of PbH_4^+ which was negative in the calculation with BPW91 functional is positive but near zero. Similarly, calculation with the UHF method also shows that all the frequencies of PbH_4^+ are positive. Irrespective of the sign of calculated frequencies, all these methods show that one vibrational mode of C_s symmetric XH_4^+ molecules has a low frequency. Also we find that the force constant corresponding to this mode is nearly zero, which implies that the H atoms in these mode are moving on nearly flat potential energy surfaces.

We see from Fig. 1 that the mirror symmetry plane of C_s structure is formed by the H-H pair and X atom. The vibrational mode with low frequency corresponds to the vibration of H-H pair about the mirror symmetry plane of C_s structure. When the H-H pair vibrates about the plane, one atom of H-H pair moves in and the other moves out of the mirror plane without change in the H-H distance. Since the force constant is nearly zero for this vibration, the motion of H-H pair is more like an internal rotation about an axis which is perpendicular to the H-H bond and passing through the X atom. This internal rotation of H-H pair in XH_4^+ is similar to the internal rotation of O-H in CH_3OH molecule[29, 30, 31]. Thus the low frequency mode in the present case does not indicate a transition state.

C. Effect of different occupations

Before proceeding further, we would like to resolve an interesting question regarding the occupation of degenerate states in charged tetrahedral XH_4 clusters. Let us ionize a charged XH_4 cluster keeping its tetrahedral structure intact. In this structure the HOMO is triply degenerate and is occupied by only 5 electrons. Hence, the HOMO of XH_4^+ is not completely filled. In such case of incompletely filled degenerate systems, there are many possible ways to fill the levels. The question is what is the right way of filling the levels and which distribution leads to the lowest energy. The usual practice in such situation is to fill all levels with fractional number of electrons, thereby keeping the symmetry intact.

To resolve the issue of occupation we have performed paramagnetic and spin-polarized calculations on tetrahedral SiH_4^+ and CH_4^+ using VASP package with local and non-local functionals for various possible electronic distributions. In paramagnetic calculations we have chosen two possible electronic distributions. The configuration with integer(2,2,1) occupation corresponds to asymmetric charge density and the other with fractional($\frac{5}{3}, \frac{5}{3}, \frac{5}{3}$) occupation corresponds to symmetric charge density. Note that the spin multiplicity of SiH_4^+ and CH_4^+ is unity. The multiplicity is preserved in charge asymmetric configuration whereas it is zero for charge symmetric configuration. In spin-polarized calculations we have chosen three possible electronic configurations. The configuration with $\frac{5}{3}\uparrow, \frac{5}{3}\uparrow, \frac{5}{3}\uparrow$ ($\frac{5}{3}\downarrow, \frac{5}{3}\downarrow, \frac{5}{3}\downarrow$) occupation corresponds to charge symmetric density and multiplicity zero. Another configuration with $1\uparrow, 1\uparrow, 1\uparrow$ ($\frac{2}{3}\downarrow, \frac{2}{3}\downarrow, \frac{2}{3}\downarrow$) occupation also corresponds to symmetric charge density but multiplicity one. The other configuration with $1\uparrow, 1\uparrow, 1\uparrow$ ($1\downarrow, 1\downarrow, 0\downarrow$) occupation corresponds to asymmetric charge density and multiplicity one. We note that in all these configurations, the Kohn-Sham wave function is a single Slater determinant. Hence the energy of the multiplet can be calculated directly using von Barth scheme[32]. The calculated total energy of tetrahedral SiH_4^+ and CH_4^+ molecules are summarized in Table. V.

From the table we see that paramagnetic calculations with local(LDA) and non-local(GGA) functionals do not give much difference between total energies of the molecules for symmetric and asymmetric charge density. On the other hand, the spin polarized calculations with local functional(LSD) favors charge symmetry for SiH_4^+ and charge asymmetry for CH_4^+ . However, for both the molecules LSD calculations show that the symmetric charge

density which has multiplicity of zero is highly unfavored. We note that in spin polarized calculations the exchange interaction is taken care of more accurately than in paramagnetic calculations. Hence the unphysical zero multiplicity is strictly not favored by spin polarized calculations. But LSD does not give consistent results for multiplicity preserved symmetric and asymmetric charge density of SiH_4^+ and CH_4^+ molecules. This is a well known shortcoming of the LSD in the context of atomic calculations[33]. This deficiency of LSD is removed by the gradient corrected functionals[34]. We have therefore done spin polarized GGA(GGA-SP) calculations on SiH_4^+ and CH_4^+ molecules. Our calculations with GGA-SP functional give the consistent result that the asymmetric charge density is the ground state configuration for both SiH_4^+ and CH_4^+ molecules in tetrahedral structure.

Apart from the total energy calculation, one more important issue is the calculation of forces in such systems. We find that enforcing charge symmetry with fractional $\frac{5}{3}, \frac{5}{3}, \frac{5}{3}$ occupation in the degenerate level of XH_4^+ in LDA calculations does not distort the tetrahedral structure. This is in contrast with the structural relaxation of tetrahedral XH_4^+ with 2,2,1 occupation of degenerate level where the molecules results in distorted structure. Although LDA does not distinguish clearly the charge symmetric and asymmetric configurations, the calculated forces are different for the two configurations. This indicates that by enforcing symmetric charge distribution, the forces on atoms of XH_4^+ molecules are incorrectly estimated. It was also noted by Janak and Williams[33] that enforcing spherical charge distribution on atoms can give rise to errors in Hellman-Feynman forces. Therefore, our results clearly indicate that the electron charge density of tetrahedral XH_4^+ is asymmetric.

D. Structural evolution

To gain insight into the mechanism of structural distortion in SiH_4^+ we look at the evolution of SiH_4^+ structure by using the VASP package with LDA functional and the conjugate gradient(CG) method[35]. We first ionize SiH_4 by holding atoms to obtain tetrahedral SiH_4^+ and then relax the structure. We find that with 64 steps the system rolls down from the initial tetrahedral to a well converged ground state structure. Each CG step corresponds to a particular geometrical configuration. The bond length between silicon atom and the distance between the hydrogen atoms as a function of CG steps are shown in Fig. 4(a). Initially one H atom moves away from the silicon atom and attains maximum Si-H bond length

of $1.92\text{\AA}$ at 8th step. Then this repelled H atom moves closer to the nearest hydrogen from 13th step which can be seen from the H-H distance shown in Fig. 4(b). The two hydrogen atoms come very close to each other to a distance of $0.84\text{\AA}$ and form an H-H pair. At this stage the structure of SiH_4^+ is C_{2v} . After 23rd step, this H-H pair gets rotated towards the other two hydrogen atoms without changing the H-H pair distance which finally results in C_s symmetric SiH_4^+ .

The summation of occupied Kohn-Sham orbital energies as a function of CG step is shown in the Fig. 4(c). From the figure we can see that the sum increases initially up to the 8th step and then decreases enormously between 13th and 23rd step. Then it fluctuates between 24 and 40th step and eventually converges to a constant value. The figure shows that the structural changes occurring from 13 and 23 are due to lowering in the orbital energies. Note that there is a minimum at 23rd step. As mentioned earlier at this CG step the structure of SiH_4^+ is C_{2v} with the H-H bond. In a sense, the sum of orbital energies shown in the figure is similar to the Walsh diagram[36] where one takes into account only the sum over orbital energies in predicting the structure. In Walsh diagram, each orbital energy is plotted as function of bond angle of a molecule. Then the minimum of the sum helps in predicting the structure. Thus in the present case the Walsh diagram would predict the symmetry of SiH_4^+ as C_{2v} symmetry and not C_s , which is the correct symmetry.

E. Role of electrostatic interaction

We now show that the distortion and formation of H-H bond can be easily understood from arguments based on electrostatic interaction between atoms of the molecule. We first try to understand the structural evolution of SiH_4^+ as discussed in the previous subsection. Let us first consider neutral SiH_4 molecule. Since H is more electronegative[37] than Si, the H atoms in SiH_4 has a small negative charge while Si will have a small positive charge. This is also shown by population analysis[18, 19, 38] according to which each hydrogen atom holds a charge of $-0.12e$ and silicon $+0.52e$. When the molecule is ionized to SiH_4^+ holding its atoms fixed, population analysis shows that the hydrogen atoms have $-0.12e$, $-0.02e$, $+0.06e$ and $+0.30e$ charge and silicon $+0.77e$ charge. Interestingly, hydrogen atoms of tetrahedral SiH_4^+ do not have equal charge although the structure is symmetric. One hydrogen atom has more positive charge than the others and gets repelled by silicon atom which is also

positively charged. This hydrogen moves away from the silicon to reduce the electrostatic repulsion and is attracted by the nearby hydrogen atom which has more electrons and forms an H-H bond. This H-H pair is positively charged and hence attracted by the nearby two hydrogen atoms which results in rotation of the pair towards the other two hydrogen atoms.

Since the electronegativities of Ge, Sn and Pb atoms are lower than that of hydrogen[37], GeH_4^+ , SnH_4^+ and PbH_4^+ have distortions and bonds similar to that of SiH_4^+ . However, as noted earlier, the structural distortion of CH_4^+ is different from that of SiH_4^+ ; two hydrogen atoms of CH_4^+ move away from each other which is in contrast with the formation of H-H bond in SiH_4^+ . We now show that the same arguments based on electronegativity and electrostatic interaction also explain the structural distortion of CH_4^+ . Since carbon has larger electronegativity than H atom, carbon in CH_4 will have negative charge while hydrogen will have positive charge. This is confirmed by population analysis which shows that a charge of -0.82e on carbon and +0.20e on each hydrogen atom. Ionizing CH_4 in the tetrahedral symmetric structure results in +0.20e, +0.23, +0.42e and +0.55e charge on H atoms and -0.40e charge on carbon. We see that the charge distribution is asymmetric and two hydrogen atoms are more positively charged than others. The two H atoms repel each other and open H-C-H bond angle as seen in Fig. 1.

Now we discuss the distortion of negatively charged molecules and explain it on the basis of the above arguments. We see from Fig. 1 that CH_4^- has the tetrahedral symmetry while other negatively charged molecules have C_{2v} symmetry. Putting an electron on CH_4 results in -1.00e on carbon and nearly zero electronic charge on H atoms. Because of this symmetrical charge distribution on hydrogen atoms, the forces on H atoms are radial and the structure of CH_4^- remain tetrahedral. In contrast, in other molecules the charge on H atoms are not equal. For example, in SiH_4^- population analysis shows a charge distribution of -0.14e, -0.15e, -0.16e and -0.37e on H atoms while silicon holds a charge of -0.18e. We can see that the H atom which is more negative gets repelled by other atoms. As a result, one of the H-Si-H bond angle opens up with the corresponding expansions in Si-H bond lengths. The trend in the electron distribution of GeH_4^- , SnH_4^- and PbH_4^- is similar to that of SiH_4^- . Therefore, the distortion of these molecules are similar to that of SiH_4^- and can be explained using the same reasoning.

F. Charge asymmetry

It is interesting to note that in most of the charged molecules the symmetry is broken. We find that the symmetry breaking is associated with the creation of asymmetry in the charge distribution on H atoms upon charging. Removing or putting an electron on these molecules without changing the tetrahedral geometry may result in charge imbalance on H atoms or equal charges on H atoms as in the case of CH_4^- . This charge imbalance or balance on H atoms depends on the electronic structure of the molecule. In particular, on the nature and occupation of the HOMO.

Using molecular orbital theory[39] we discuss the nature of HOMO in these molecules and show how it contributes to charge distribution among H atoms. In Fig. 5, we show the MO construction of XH_4 from linear combinations of atomic orbitals. As shown in the figure, there are eight MOs, the lower four are bonding states and the upper four are antibonding states. The bonding and antibonding states have either A_1 symmetry or T_2 symmetry. A_1 level is non-degenerate while T_2 level is triply degenerate. While the bonding energy level of A_1 symmetry is always lower in energy than T_2 , the order of antibonding levels depends on the molecule. For CH_4 antibonding A_1 level is lower in energy than T_2 while for SiH_4 , GeH_4 , SnH_4 and PbH_4 antibonding T_2 level is lower than A_1 . From the figure we can see that the MO with A_1 symmetry is a linear combination of valence atomic s orbitals of X and H atoms. The MO with T_2 symmetry is a linear combination of valence p orbitals of X and s orbitals of H atoms. Because of mixing between p and s character, T_2 orbitals have strong directionality along H atoms. If T_2 orbital is fully occupied, it has equal weight along H atoms and results in a uniform charge distribution on H atoms as in the case of neutral XH_4 molecules. If T_2 orbital is incompletely filled with unequal electrons, it has unequal weight along H atoms and results in a non-uniform charge distribution on H atoms.

Removing an electron from tetrahedral XH_4 results with 5 electrons and the T_2 orbital is incompletely filled. Hence the charges on H atoms of tetrahedral XH_4^+ molecules are not equal. This can be seen from Fig. 6(a) and (b) which show the charge densities of SiH_4 and SiH_4^+ in tetrahedral structure. We see from the figure that while the charge densities on H atoms of SiH_4 are same, the charge densities on H atoms of SiH_4^+ are different. In the case of negatively charged molecules the HOMO can be of T_2 or A_1 symmetry. The HOMO of CH_4^- has A_1 symmetry. Since A_1 is a linear combination of valence s atomic orbitals with

equal weights from hydrogen atoms, the charge distribution is equal on H atoms of CH_4^- as shown in Fig. 6(c). The higher electron density around the carbon atom as seen in the figure can be attributed to the higher electronegativity of carbon compared to hydrogen. The HOMO of other negatively charged molecules has T_2 symmetry and is singly occupied. This leads to asymmetric charge distribution on H atoms. Note that since A_1 level in CH_4^- is non-degenerate, CH_4^- does not show the Jahn-Teller effect, and also no charge asymmetry. On the other hand, if the HOMO has T_2 symmetry and is not completely occupied as in the case of SiH_4^+ , the molecule will show the Jahn-Teller effect and the charge asymmetry. This implies that the creation of the charge asymmetry and the Jahn-Teller effect are intimately connected.

IV. CONCLUSIONS

We have investigated the ground state structures of neutral and charged XH_4 molecules using first principle electronic structure methods. We find that charging XH_4 molecules can break the symmetry and change their structures drastically. Negatively charged molecules are distorted by pushing two hydrogen atoms away from each other while the positively charged molecules get distorted with an unusual H-H bond formation except for $\text{X}=\text{C}$. The calculations done by various methods and the charge density analysis confirm the existence of H-H bond. Furthermore, vibrational analysis shows that the strength of H-H bond in the positively charged molecule is stronger than H_2^+ and weaker than H_2 . We find that one vibrational mode of C_s symmetric XH_4^+ molecules has low frequency and corresponds to internal rotation of the H-H pair. Our vibrational analysis indicates that the molecules are not transition states. The fragmentation behavior of XH_4^+ into XH_2 and H_2 shows that the stability decreases from CH_4^+ to PbH_4^+ .

From the total energy calculations using LDA, LSD, GGA and GGA-SP functionals we have shown that the ground state electronic structure of a charged tetrahedral molecule has asymmetric charge density corresponding to integer occupation. We have provided an insight into the symmetry breaking and shown that the distortion and unusual H-H bond formation can be understood easily using simple arguments based on charge asymmetry and electrostatic interaction between the atoms. Unequal charges on hydrogen atoms give rise to different electrostatic forces on different atoms and lead to distortion. On the other hand

if the charges are equal on hydrogen atoms, the structural symmetry of the molecule will be preserved. We have shown how the charge asymmetry arises due to character of the HOMO and eventually leads to the symmetry breaking.

Acknowledgments

It is a pleasure to thank Drs. R. Ramaswamy, D. D. Sarma, S. C. Agarwal and N. Sathya-murthy for helpful discussions and comments. This work was supported by the Department of Science and Technology, New Delhi via project No. SP/S2/M-51/96.

-
- [1] I. B. Bersuker, Chem. Rev. **101**, 1067 (2001).
 - [2] M. D. Sturge in *The Jahn-Teller effect in solids* edited by F. Sietz, D. Turnbull and H. Ehrenreich, volume 20 of Solid State Physics, Advances in Research and Applications (1967) p. 91.
 - [3] R. Englman, *The Jahn-Teller effect in molecules and crystals* (Wiley-Interscience, New York, 1972).
 - [4] R. F. Frey and E. R. Davidson, J. Chem. Phys. **89**, 4227 (1988).
 - [5] F. D. Proft and P. Geerlings, Chem. Phys. Lett. **262**, 782 (1996).
 - [6] T. Kudo and S. Nagase, Chem. Phys. **122**, 233(1988).
 - [7] R. Caballol, J. A. Catala and J. M. Problet, Chem. Phys. Lett. **130**, 278 (1986).
 - [8] A. R. Porter, O. K. Al-Mushadani, M. D. Towler and R. J. Needs, J. Chem. Phys. **114**, 7795 (2001).
 - [9] S. D. Wetmore, R. J. Boyd, L. A. Eriksson and A. Laaksonen, **110**, J. Chem. Phys, 12059 (1999).
 - [10] Z. Vager, E. P. Kanter, G. Both, P. J. Cooney, A. Faibis, W. Koenig, B. J. Zabransky and D. Zajfman, Phys. Rev. Lett. **57**, 2793, (1986).
 - [11] R. Signorell and F. Merkt, J. Chem. Phys. **110**, 2309(1999).
 - [12] J. M. G. De La Vega, A. Segovia, J. F. Rico and J. I. Fernández-Alonso, J. Mol. Struct. **123**, 203(1985).
 - [13] R. J. Boyd, K. V. Darvesh and P. D. Fricker, J. Chem. Phys. **94**, 8083 (1991).

- [14] R. Signorell and M. Sommovilla, J. Electron Spectroscopy and Related Phenomena, **108**, 169(2000).
- [15] K. Takeshita, J. Chem. Phys. **86**, 329(1987).
- [16] R. F. Frey and E. R. Davidson, J. Chem. Phys. **88**, 1775(1988).
- [17] G. Kresse and J. Furthmüller, Comput. Mater.Sci. **6** 15 (1996); Phys. Rev. **B 54**, 11169 (1996).
- [18] Gaussian 98, Revision A.11.1, M. J. Frisch *et al* Gaussian, Inc., Pittsburgh PA, 2001.
- [19] J. B. Foresman and A. Frisch, *Exploring Chemistry with Electronic Structure Methods* (Second Edition, Gaussian, Inc. Pittsburgh, PA, 1993).
- [20] D. M. Ceperley and B. J. Alder, Phys. Rev. Lett. **45**, 566 (1980).
- [21] J. P. Perdew and A. Zunger, Phys. Rev. **B 23**, 5048 (1981).
- [22] J. Neugebauer and M. Scheffler, Phys. Rev. **B 46**, 16067 (1992).
- [23] G. Makov and M. C. Payne, Phys. Rev. **B 51**, 4014(1995), G. Makov, R. Shah and M. C. Payne, Phys. Rev. **B 53**, 15513(1996).
- [24] D. Marx and M. Parrinello, Science, **284**, 59(1999).
- [25] D. Marx and A. Savin, Angew. Chem. Int. Ed. Engl. **36**, 2077(1997).
- [26] D. W. Boo and Y. T. Lee, J. Chem. Phys, **103**, 514(1995).
- [27] D. Balamurugan and R. Prasad, Phys. Rev. **B 64**, 205406(2001).
- [28] A. M. Coats, D. C. McKean and D. Steele, J. Mol. Struct. **320**, 269(1994).
- [29] P. Y. Ayala and H. B. Schlegel, J. Chem. Phys. **108**, 2314(1998).
- [30] C. R. Quade, J. Mol. Spect. **188**, 190(1998).
- [31] Y. Duan, R. Wang and I. Mukhopadhyay, Chem. Phys. **280**, 119(2002).
- [32] U. von Barth, Phys. Rev. A **20**, 1693(1979).
- [33] J. F. Janak and A. R. Williams, Phys. Rev. B **23**, 6301(1981).
- [34] F. W. Kutzler and G. S. Painter, Phys. Rev. Lett. **59**, 1285(1987).
- [35] We have also looked at the evolution of SiH_4^+ structure using the steepest descent method. Although the convergence is slow, the results are similar to those shown in Fig. 4.
- [36] See, for example, I. N. Levine, *Quantum Chemistry* (Prentice-Hall, Englewood Cliffs, New Jersey, 1991) p. 503-506.
- [37] J. Robles and L. J. Bartolotti, J. Am. Chem. Soc. **106**, 3723(1984).
- [38] The numbers shown in the text are obtained from the natural population analysis using

BPW91 exchange correlation potential. The numbers obtained by the Mulliken analysis are somewhat different but show the same trend and thus do not change our conclusions.

- [39] See, for example, A. A. Levin, *Solid State Quantum Chemistry* (McGraw Hill Book Company, New York, 1977) p. 34.

FIGURE CAPTIONS

Fig. 1. Ground state structures of neutral and charged XH_4 molecules.

Fig. 2 Electron density contour of SiH_4^+ in the plane of Si and the two bonded H atoms.

Fig. 3 The total energy of SiH_4^+ , H_2 and H_2^+ molecules as a function of H-H distance. The total energy of the molecule is given with respect its ground state energy.

Fig. 4. (a) The Si-H bond lengths and (b) H-H distances in SiH_4^+ as a function of CG steps. The H atoms have the same label as those of SiH_4^+ in Fig. 1. (c) The sum of energy levels of SiH_4^+ as a function of CG steps.

Fig. 5. A schematic energy-level diagram showing the construction of molecular orbitals of XH_4 molecule from atomic orbitals. The levels on the extreme left side are the valance states of X atoms and the level on the extreme right is the valance state of H atom. The levels which are in the center are the molecular states of XH_4 .

Fig. 6. The isosurface contour of valance electrons of (a) SiH_4 (b) SiH_4^+ and (c) CH_4^- obtained by using the Gaussian package. The value of the contour is chosen such that one able to distinguish the difference between the charge density on the H atoms. The densities shown in figure have the contour value of 0.20 for SiH_4 and SiH_4^+ molecules and 0.3 for CH_4^- .

Table. I. Relative energies (in eV) of XH_4^+ molecules for various symmetries with respect to the ground state structures. The results are obtained using the Gaussian98 with the BPW91 exchange-correlation functional.

	C_{2v}	D_{2d}	C_{3v}	C_s
CH_4^+	0.11	0	-	-
SiH_4^+	-	0.73	0.73	0
GeH_4^+	0.83	0.94	0.87	0
SnH_4^+	1.36	1.46	1.35	0
PbH_4^+	1.80	1.91	1.72	0

Table. II Properties of H-H bond calculated using the Gaussian98 package with the BPW91 exchange-correlation functional.

	SiH_4^+	GeH_4^+	SnH_4^+	PbH_4^+	H_2	H_2^+
H-H bond length($\text{\AA}$)	0.80	0.78	0.77	0.76	0.75	1.12
H-H frequency (cm^{-1})	3598	3911	4061	4134	4349	1976
H-H force constant($\text{eV}/\text{\AA}$)	7.69	9.08	9.79	10.15	11.23	2.32

Table. III. The energy required to fragment XH_4^+ into XH_2^+ and H_2 . The results are obtained using the Gaussian98 package with the BPW91 exchange-correlation functional.

Reactant	product 1	product 2	Fragmentation energy(eV)
CH_4^+	CH_2^+	H_2	3.03
SiH_4^+	SiH_2^+	H_2	0.69
GeH_4^+	GeH_2^+	H_2	0.42
SnH_4^+	SnH_2^+	H_2	0.28
PbH_4^+	PbH_2^+	H_2	0.21

Table. IV The calculated vibrational frequencies ν (in cm^{-1}) and force constant k (in $\text{eV}/\text{\AA}$) of neutral and charged XH_4 molecules using the Gaussian98 package with the BPW91 exchange-correlation potential.















	Neutral molecules									
	CH_4		SiH_4		GeH_4		SnH_4		PbH_4	
	ν	k	ν	k	ν	k	ν	k	ν	k
1	1298	1.17	879	0.50	788	0.38	676	0.28	644	0.25
2	1298	1.17	880	0.50	788	0.38	676	0.28	644	0.25
3	1298	1.17	881	0.50	788	0.38	676	0.28	644	0.25
4	1521	1.37	949	0.53	901	0.48	735	0.32	703	0.29
5	1521	1.37	950	0.54	901	0.48	735	0.32	703	0.29
6	2975	5.26	2175	2.81	2058	2.51	1856	2.05	1769	1.86
7	3090	6.19	2191	2.97	2071	2.59	1856	2.07	1786	1.91
8	3090	6.19	2191	2.97	2072	2.59	1857	2.07	1786	1.91
9	3090	6.19	2191	2.97	2072	2.59	1857	2.07	1787	1.91
	Positively charged molecules									
	CH_4^+		SiH_4^+		GeH_4^+		SnH_4^+		PbH_4^+	
	ν	k	ν	k	ν	k	ν	k	ν	k
1	470	0.15	256	0.04	136	0.01	46	0.00	-104	0.01
2	471	0.15	616	0.25	510	0.16	380	0.09	316	0.06
3	1022	0.78	680	0.28	532	0.17	394	0.09	339	0.07
4	1253	0.93	766	0.35	542	0.18	430	0.11	396	0.09
5	1420	1.20	850	0.45	658	0.26	495	0.15	425	0.11
6	2642	4.26	1077	0.69	781	0.37	666	0.27	631	0.24
7	2758	4.52	2164	2.83	1957	2.29	1729	1.78	1506	1.35
8	2851	5.32	2242	3.15	2046	2.54	1807	1.96	1623	1.57
9	2852	5.32	3598	7.69	3911	9.08	4061	9.79	4134	10.15


TABLE IV: Continued


	Negatively charged molecules									
	CH_4^-		SiH_4^-		GeH_4^-		SnH_4^-		PbH_4^-	
	ν	k	ν	k	ν	k	ν	k	ν	k
1	1133	0.88	690	0.30	614	0.23	491	0.15	347	0.07
2	1133	0.88	759	0.36	729	0.33	619	0.23	469	0.13
3	1133	0.88	790	0.38	768	0.35	645	0.25	673	0.27
4	1372	1.12	955	0.58	884	0.48	784	0.37	675	0.27
5	1372	1.12	1000	0.59	934	0.52	794	0.37	692	0.29
6	2537	3.82	1323	1.09	1099	0.73	1067	0.68	766	0.35
7	2631	4.55	1429	1.22	1323	1.04	1245	0.92	1124	0.75
8	2631	4.55	1958	2.32	1786	1.91	1607	1.54	1472	1.29
9	2631	4.55	1962	2.37	1804	1.96	1615	1.56	1489	1.32


Table. V Relative energies(in eV) of SiH_4^+ and CH_4^+ in tetrahedral structure for various electronic configuration with respect to charge asymmetric configuration, using different exchange-correlation functionals.


	configuration	SiH_4^+	CH_4^+
LDA	2, 2, 1	0.000	0.000
	$\frac{5}{3}, \frac{5}{3}, \frac{5}{3}$	0.001	0.001
GGA	2, 2, 1	0.000	0.000
	$\frac{5}{3}, \frac{5}{3}, \frac{5}{3}$	0.030	0.000
LSD	$1\uparrow, 1\uparrow, 1\uparrow(1\downarrow, 1\downarrow, 0\downarrow)$	0.000	0.000
	$\frac{5}{3}\uparrow, \frac{5}{3}\uparrow, \frac{5}{3}\uparrow(\frac{5}{3}\downarrow, \frac{5}{3}\downarrow, \frac{5}{3}\downarrow)$	0.139	0.296
	$1\uparrow, 1\uparrow, 1\uparrow(\frac{2}{3}\downarrow, \frac{2}{3}\downarrow, \frac{2}{3}\downarrow)$	-0.028	0.036
GGA-SP	$1\uparrow, 1\uparrow, 1\uparrow(1\downarrow, 1\downarrow, 0\downarrow)$	0.000	0.000
	$\frac{5}{3}\uparrow, \frac{5}{3}\uparrow, \frac{5}{3}\uparrow(\frac{5}{3}\downarrow, \frac{5}{3}\downarrow, \frac{5}{3}\downarrow)$	0.215	0.333
	$1\uparrow, 1\uparrow, 1\uparrow(\frac{2}{3}\downarrow, \frac{2}{3}\downarrow, \frac{2}{3}\downarrow)$	0.070	0.105
















(a)



(b)



(c)